

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/00	A2	(11) International Publication Number: WO 98/51788 (43) International Publication Date: 19 November 1998 (19.11.98)
(21) International Application Number: PCT/NL98/00259 (22) International Filing Date: 11 May 1998 (11.05.98) (30) Priority Data: 97201423.7 12 May 1997 (12.05.97) EP (34) Countries for which the regional or international application was filed: NL et al. (71) Applicant (for all designated States except US): NEDERLANDSE ORGANISATIE VOOR TOEGEPAST-NATUURWETENSCHAPPELIJK ON- DERZOEK TNO [NL/NL]; Schoemakerstraat 97, NL-2628 VK Delft (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): QUAX, Paulus, Hubertus, Andreas [NL/NL]; Jacob van Heemskercklaan 59, NL-2253 JX Voorschoten (NL). VERHEIJEN, Johan, Hendrikus [NL/NL]; Verdijkstraat 14, NL-2651 VB Berkel en Rodenrijs (NL). (74) Agent: SMULDERS, Th., A., H., J.; Vereenigde Octrooibu- reaux, Nieuwe Parklaan 97, NL-2587 BN The Hague (NL).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>
(54) Title: METHOD AND CONSTRUCT FOR INHIBITION OF CELL MIGRATION (57) Abstract A recombinant nucleic acid molecule comprising a vector useful for transfection or transduction of mammalian cells, wherein said vector contains a nucleic acid insertion encoding an expressible hybrid polypeptide or protein which comprises a domain with a binding function and a domain with an effector function. The domain with a binding function may comprise a receptor binding domain, and the domain with an effector function may have enzymatic activity, in particular protease inhibitor activity. The vector may be a viral (e.g. adenovirus or retrovirus) or non-viral vector useful for transfection or transduction of mammalian cells. The nucleic acid insertion encoding an expressible hybrid polypeptide or protein may be under the control of a cell- or tissue-specific promoter. A process for preventing local proteolytic activity, extracellular matrix degradation, cell migration, cell invasion, or tissue remodeling, comprising transfecting or transducing the cells involved or cells in their environment with the recombinant nucleic acid molecule to obtain local expression of the hybrid polypeptide or protein encoded thereby. A process for producing the hybrid polypeptide or protein by transfecting or transducing mammalian cells with the recombinant nucleic acid molecule to obtain expression and optionally recovering the hybrid polypeptide or protein produced.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

Title: Method and Construct for inhibition of cell migration

FIELD OF THE INVENTION

The invention is in the field of therapeutic means and therapeutic methods for treatment of diseases in which cell migration and/or tissue remodeling occurs. Furthermore, the invention is in the field of biotechnology, in particular recombinant DNA technology and gene therapy.

BACKGROUND OF THE INVENTION

Migration of cells is an essential step in many physiological and pathological processes in which tissue remodeling occurs, such as tumor metastasis, wound healing, restenosis, angiogenesis or rheumatic arthritis. Migrating cells have to pass through the surrounding extracellular matrix. Limited proteolytic degradation of the components of the extracellular matrix is often seen during cell migration. To mediate this cell migration migrating cells produce, or recruit from their direct environment, proteolytic enzymes, such as plasminogen activators, metalloproteinases or elastases. Induction of cell migration e.g. during tumor metastasis or wound healing often correlates with the induction of the production of these enzymes.

Although the involvement of proteolytic enzymes in cell migration under pathophysiological conditions is well accepted, little attempts have been made to inhibit cell migration by inhibiting these proteolytic enzymes. A possible explanation for the limited use of protease inhibitors is the fact that these proteolytic enzymes are involved in many processes both pathological and physiological (including fibrinolysis, wound healing, growth factor activation etc.) and that inhibition of these protease systems by systemically applied protease inhibitors might have either strong side effects or may lead to a diffusion or clearance of the inhibitory compounds without having a strong effect on the local cell migration processes.

Another problem in the use of protease inhibitors to interfere in cell migration and tissue remodeling is that proteases mediating these processes can bind to receptors at the cell surface. In this way the proteolytic enzymes might
5 be active locally in a pericellular microenvironment where they are protected against the action of the present inhibitors.

It has been disclosed that conjugates between the receptor binding part of u-PA (the aminoterminal fragment or
10 ATF) and urinary trypsin inhibitor produced in vitro, inhibit migration of tumor cells in vitro (Kobayashi, Gotoh, Hirashima, Fujie, Sugino and Terao, Inhibitory effect of a conjugate between human urokinase and urinary trypsin inhibitor on tumor cell invasion in vitro. J. Biol. Chem.
15 (1995) 270, 8361-8366). The conjugate these authors have used is made synthetically by mixing the isolated ATF fragments with the trypsin inhibitor.

Recently it has been disclosed that these conjugates also can be produced recombinantly (WO 97/25422).

20 A comparable construct consisting of a receptor binding u-PA fragment and its inhibitor PAI-2, to be produced recombinantly in yeast, has been described to inhibit tumor cell migration in WO 92/02553 (PCT/GB91/01322). In this way they have made a protease inhibitor that can bind to a
25 specific receptor at the cell surface, the urokinase receptor, and this inhibitor can inhibit cell migration (in vitro). As to the use of these constructs in vivo, a problem is the application to and the prolonged presence at the site of desired action in vivo.

30

SUMMARY OF THE INVENTION

This invention provides a recombinant nucleic acid molecule comprising a vector useful for transfection or transduction of mammalian, e.g. human, cells, wherein said
35 vector contains a nucleic acid insertion encoding an expressible hybrid polypeptide or protein which comprises a

domain with a binding function and a domain with an effector function. Herein, the domain with a binding function preferably comprises a receptor binding domain, and the domain with an effector function preferably has enzymatic activity, most preferably protease inhibitor activity.

Preferably, the receptor binding domain is selected from the group consisting of urokinase receptor binding domain of urokinase, receptor binding domain of epidermal growth factor, receptor associated protein that binds to LDL Receptor related protein (α_2 -macroglobulin receptor) and VLDL Receptor.

Preferably, the domain with an effector function has protease inhibitor activity and comprises a protease inhibitor or active part thereof, said protease inhibitor being selected from the group consisting of (bovine) pancreatic trypsin inhibitor, (bovine) splenic trypsin inhibitor, urinary trypsin inhibitor, tissue inhibitor of matrix metalloproteinase 1, tissue inhibitor of matrix metalloproteinase 2, tissue inhibitor of matrix metalloproteinase 3, and elastase inhibitor. The domain with an effector function may comprise (an active part of) two or more different protease inhibitors, or two or more copies of (an active part of) a protease inhibitor, or both.

Preferably, the vector is selected from the group consisting of viral and non-viral vectors useful for transfection or transduction of mammalian cells. The vector may be an adenovirus vector or a retrovirus vector useful for transfection or transduction of human cells.

The nucleic acid insertion encoding an expressible hybrid polypeptide or protein may be under the control of a cell- or tissue-specific promoter, such as an endothelial cell-specific promoter, or a vascular smooth muscle cell-specific promoter, or a liver-specific promoter.

This invention furthermore provides a process for preventing local proteolytic activity, extracellular matrix degradation, cell migration, cell invasion, or tissue

remodeling, comprising transfecting or transducing the cells involved or cells in their environment with a recombinant nucleic acid molecule as defined herein to obtain local expression of the hybrid polypeptide or protein encoded by
5 said nucleic acid molecule.

Also, this invention provides a process for producing a hybrid polypeptide or protein which comprises a domain with a binding function and a domain with an effector function, comprising transfecting or transducing mammalian
10 cells with a recombinant nucleic acid molecule as defined herein to obtain expression of the hybrid polypeptide or protein encoded by said nucleic acid molecule, and optionally recovering the hybrid polypeptide or protein produced.

15 BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 schematically depicts the plasmids pCRII-uPA (left) and pCRII-ATF (right).

Figure 2 schematically depicts the plasmid pCRII-ATF-BPTI.

20 Figure 3 schematically depicts the plasmid pMAD5-ATF-BPTI.

Figure 4 shows the results of proteolytic matrix degradation experiments.

25 DETAILED DESCRIPTION OF THE INVENTION

The present invention relates to the use of hybrid proteins in which a receptor binding domain is linked to a functional protein in order to induce a local action of this protein and to prevent systemic effects and/or diffusion. In
30 particular this invention relates to such hybrid proteins that might be produced by a subset of cells as target cells after transfection or transduction with expression vectors. More specifically the invention relates to the use of such expression vectors, coding for hybrid proteins consisting of
35 a receptor binding domain and a protease inhibitor domain, for the prevention of cell migration and tissue remodeling by

inhibition of proteases at the surface of migrating or invading cells.

The method and construct described in the present invention can be applied as therapy in diseases in which cell migration and/or tissue remodeling occurs.

The present invention addresses the solution of several negative aspects involved in the above described use of inhibitors according to the prior art:

- High local concentrations of hybrid proteins in the direct environment of the target cells can be obtained by production of the protein by the migrating cells themselves or cells in their immediate environment. This production can be induced by transfection or transduction of a certain subset of the cell population with a suitable vector encoding the hybrid protein. For this purpose, one may use recombinant adenoviral vectors, retroviral vectors, plasmid vectors, etc.
- Diffusion of the inhibitor and systemic side effects are prevented by binding the hybrid protein (by its receptor binding domain) to the cell surface of the target cell. Local expression of this hybrid protein also contributes to the reduction of systemic side effects, while the negative effect of diffusion of the protein is reduced by the production at the site where action is required. The local expression of the hybrid protein in specific subpopulations of cells, e.g. endothelial cells prone to migrate during angiogenesis, can be enhanced using cell type-specific or tissue-specific expression vectors, in which the expression of the protein is under control of a promoter with cell type-specific or tissue-specific regulatory elements.
- Binding of a protease inhibitor to a cell surface receptor can locate the inhibitor close to its molecular target, the cell surface bound proteolytic enzyme. Local inhibition of the proteolytic activity in the pericellular microenvironment may be achieved in this way.
- Binding of a protease inhibitor to a cell surface receptor for a proteolytic enzyme, such as the urokinase

receptor, may have an additional inhibitory effect. It prevents the binding of the proteolytic enzyme to its receptor, and thus strongly reduces the action of this enzyme as has been shown for blocking the binding of u-PA to its
5 receptor which can strongly inhibit cell migration.

Hybrid proteins, for which the expression vectors (e.g. adenoviral or retroviral expression vectors) contain the encoding DNA sequences, might contain a region that binds to a cell surface receptor and that is not subsequently
10 internalized. Receptor binding domains that can be used for this purpose are e.g. the u-PAR binding domain of urokinase plasminogen activator, the receptor binding domain of epidermal growth factor, the receptor associated protein (RAP) that binds to the LDL-R related protein (LRP), also
15 called α_2 -macroglobulin receptor, and the VLDL-receptor.

The inhibitor part of the encoded hybrid protein might consist of various protease inhibitors such as bovine pancreatic trypsin inhibitor, also called aprotinin or Trasylol[®], other trypsin inhibitors such as urinary trypsin
20 inhibitor, inhibitors for matrix-degrading metalloproteinases such the tissue inhibitors of metalloproteinases TIMP-1, TIMP-2 and TIMP-3, or variants thereof. Also inhibitors for other proteases like elastase are very suitable for being incorporated into the expression vector containing the DNA
25 sequences encoding the hybrid proteins. Multiple copies of the DNA sequences encoding the functional protein part of the hybrid protein e.g. the inhibitor part, or combinations of different inhibitors or derivatives thereof might be incorporated into the DNA construct in the expression vector.

30 Another very attractive possibility would be to use such an expression vector encoding hybrid receptor binding protein to apply any functional protein that should exert its action in the local environment of the target cell, e.g. a protease involved in the activation of a growth factor or an
35 other e.g. vasoregulatory component.

The action of the functional protein or protein domains of the hybrid protein is localized to the direct microenvironment of the target cells by binding of the receptor binding domain to a receptor at the surface of the target cells. Production of the hybrid protein in the direct environment of the target cells or even by the target cells themselves can be obtained by transfection or transduction of these cells by the use of expression vectors that might be based on a non-viral or an adeno- or retroviral vector system. Expression in specific cell or tissue types might be achieved by the use of specific promoter elements in the expression vectors. For example, for endothelial cell-specific expression (elements of) the promoter region of the human or murine pre-pro-endothelin gene (HUMEDN1B and MMU07982, respectively, GENBANK) can be used, for vascular smooth muscle cell-specific expression (elements of) the promoter region of the human vascular smooth muscle α -actin gene (HUMACTSA, GENBANK) can be used, and for liver-specific expression the promoter of the human albumin gene (HUMALBGC, GENBANK) can be used.

Local delivery of these vectors might be obtained using various commonly used methods, including catheters, topically applied gels containing the vectors or targeted delivery systems. For site-specific delivery to the vessel wall, e.g. to prevent restenosis and vessel wall remodeling after angioplasty, special catheters can be used. At the moment double balloon catheters, channeled balloon catheters, multiple needle catheters and balloon catheters coated with a vector containing a hydrogel are being used for vessel wall-specific delivery. Other ways to deliver the vectors directly into the vessel wall are the use of stents coated with vector containing coatings, topical application of vector containing hydrogels to the outside of the blood vessel or ex vivo delivery directly into the blood vessel during transplantation surgery. Ex vivo transduction of proliferating cells using retroviral vectors followed by a reinjection may

also be used to deliver the vector constructs at the site where their action is required.

The present application will be described herein-
5 after in further detail, while referring to the following examples. It is to be noted that these examples merely serve to illustrate the invention, not to restrict it.

EXAMPLE 1

10 An expression plasmid encoding the aminoterminal fragment of urokinase plasminogen activator (u-PA), amino acids 1-138, hereafter referred to as ATF, can be constructed by deleting the DNA sequences encoding amino acids 139 till 401 in an expression plasmid for the full length u-PA using a
15 polymerase chain reaction (PCR) with the following oligonucleotides: 5'-cccgggcttttttccatctgcgcagtc-3' and 5'-agggtcaccaaggaagagaatggc-3'. After amplification by PCR the newly formed DNA fragment can be circularized by ligation to restore the circular character of the expression plasmid.
20 In this way an expression plasmid encoding the ATF and the C terminal last 11 amino acid residues including the stop codon can be constructed.

The sequence of the thus formed DNA construct encoding the u-PA ATF fragment then is determined and compared
25 with the predicted sequence as a control for possible mutations introduced during the construction procedure.

The construction pCRII-ATF from pCRII-uPA using PCR is shown in Figure 1. In figure 1, the area indicated between the lines was removed during the PCR amplification, resulting
30 in the ATF plasmid. The plasmid pCRII-uPA is shown to the left, plasmid pCRII-ATF to the right.

EXAMPLE 2

DNA fragments encoding amino acid residues 36-93 of
35 bovine pancreatic trypsin inhibitor (BPTI) and the homologous amino acid residues of bovine spleen trypsin inhibitor (BSTI)

can be isolated by performing a PCR reaction on genomic DNA isolated for bovine aortic endothelial cells using the following oligonucleotides: 5'-*tcgcgc*acctgacttctgcctagagc-3' covering nucleotides 2509 to 2533 (with modifications, indicated in *italics*, in the 5' region of the oligonucleotide to introduce a NruI site (underlined) for cloning purposes) of the BPTI gene according to the published sequence (GENBANK, BTBPTIG), and nucleotide 2442 to 2462 of the BSTI gene according to the published sequence (GENBANK, BTBSTIG) and 5'-*ggtcacc*cagggcccaatattaccacc-3' covering nucleotides 2677 to 2704 of the BPTI gene and 2610 to 2636 of the BSTI gene (modified in the indicated nucleotides (*italics*) to introduce a BstEII and a SspI site respectively (underlined)). The amplified DNA fragments then were cloned into an appropriate plasmid vector, pCRII or pUC13, and then the exact sequence of the amplified DNA fragments in the isolated clones was analyzed to differentiate between BPTI and BSTI which have a very high degree of homology.

20 EXAMPLE 3

The DNA fragment encoding amino acids 1 to 207 of the human tissue inhibitor of metalloproteinase type 1 is isolated by performing a reverse transcriptase polymerase chain reaction on total RNA isolated from human foreskin fibroblasts by using the following oligonucleotides 5'-agagagacaccagagaacccaccat-3' covering nucleotides 41 to 65 of the human TIMP-1 cDNA (according to the sequence in GENBANK HSTIMPR) and 5'-tcattgtccggaagaaagatgggag-3' covering nucleotides 740 till 755. The amplified DNA fragment was cloned into an appropriate host vector, pUC13, and then the exact sequence of the amplified DNA fragment in the isolated clones was analyzed.

EXAMPLE 4

35 For construction of a recombinant adenovirus containing sequences encoding the ATF.BPTI hybrid protein,

this sequence is cloned in the adenoviral vector construction adapter and expression plasmid pMAD5. This plasmid contains part of the wildtype adenovirus type 5 DNA sequences, a Major Late Promoter (MLP), and a poly-adenylation (polyA) signal and can be used as either an expression vector or a shuttle vector to construct a recombinant adenovirus. This plasmid was derived from plasmid pMLP10 as follows. First pMLP10-lin was constructed by insertion of a synthetic DNA fragment with unique sites for the restriction endonucleases MluI, SplI, SnaBI, SpeI, AsuII and MunI into the HindIII site of pMLP10. Subsequently, the adenovirus BglII fragment spanning nt 3328 to 8914 of the Ad5 genome was inserted into the MunI site of pMLP10-lin. Finally, the SalI-BamHI fragment was deleted to inactivate the tetracycline resistance gene, resulting in plasmid pMAD5. To clone the ATF.BPTI sequence into the pMAD5 plasmid between the MLP promoter and the polyA signal the following strategy has been followed.

Starting from a pCRII plasmid in which a 1373 base pair fragment of the uPA cDNA was cloned, a PCR reaction with the oligonucleotides 5'-ccccgggcttttttccatctgcgagtc-3' (SmaI site underlined and nucleotides changed in *italics*) and 5'-agggtccaccaaggaagagaatggc-3' (BstEII site underlined and nucleotides changed in *italics*) was performed as described in example 1 to make a pCRII-ATF plasmid (see figure 1). Subsequently this pCRII-ATF plasmid was digested with the restriction enzymes SmaI and BsteII. In parallel the pCRII-BPTI plasmid was digested with the restriction enzymes NruI and BsteII and the BPTI containing fragment was cloned into the pCRII-ATF plasmid (see figure 2). The construction pCRII-ATF-BPTI is shown in Fig. 2.

In a next step the ATF-BPTI sequence was cloned into pMAD5. This was done by digestion of the pCRII-ATF-BPTI plasmid with the restriction enzymes EcoRV and SpeI, isolation of the ATF-BPTI encoding DNA fragment and cloning of this fragment into the SnaBI and SpeI digested pMAD5

plasmid. The cloning was tested by restriction analysis and sequence analysis.

The pMAD5-ATF-BPTI shuttle vector for the construction of ATF-BPTI adenoviral vector is shown in Figure 3.

5

EXAMPLE 5

In a similar way as described in example 4 for pMAD5-ATF-BPTI a plasmid containing the BSTI-gene (pMAD5-ATF-BSTI) was constructed using the pCRII-BSTI plasmid instead of
10 the pCRII-BPTI plasmid.

EXAMPLE 6

For construction of a recombinant adenovirus containing sequences encoding the ATF-TIMP1 hybrid protein,
15 this sequence is cloned in the pMAD5 expression plasmid. This plasmid contains part of the wildtype adenovirus type 5 DNA sequences, a Major Late Promoter (MLP), and a polyadenylation (polyA) signal and can be used as either an expression vector or a shuttle vector to construct a recombinant adenovirus. To
20 clone the ATF-TIMP1 sequence into the pMAD5 plasmid between the MLP promoter and the polyA signal, the following strategy has been followed.

Starting from a pCRII plasmid in which a 1373 base pair fragment of the uPA cDNA was cloned, a PCR reaction with
25 the oligonucleotides 5'-cccgggcttttttccatctgcgagtc-3' and 5'-agggtcaccaaggaagagaatggc-3' was performed as described in example 1 to make a pCRII-ATF plasmid (see figure 1). Subsequently this pCRII-ATF plasmid was digested with the restriction enzymes SmaI and BsteII.

30 In parallel a fragment of the cDNA of TIMP1 in pUC13-TIMP1 encoding amino acid residues 1 to 184 of the mature protein, but lacking the signal peptide and the stop codon, was amplified using the following oligonucleotides 5'-tcgcgatgcacctgtgtcccacc-3' and
35 5'-ggtcacccaaatattggctatgtgggaccgcaggg-3'. These oligonucleotides contain recognition sites for the restriction

enzymes NruI (first oligonucleotide, underlined) and BstEII and SspI respectively (second oligonucleotide, underlined); these sites are needed for the cloning procedure.

The amplified DNA fragment was cloned into a pCRII
5 vector and called pCRII-TIMP1. This vector was subsequently digested with the restriction enzymes NruI and BstEII and the TIMP1 containing DNA fragment was cloned into the pCRII-ATF plasmid (see figure 1).

In a next step the ATF-TIMP sequence was cloned
10 into pMAD5. This was done by digestion of the pCRII-ATF-TIMP plasmid with the restriction enzymes EcoRV and SpeI, isolation of the ATF-TIMP encoding DNA fragment and cloning of this fragment into the SnaBI and SpeI digested pMAD5 plasmid. The cloning was tested by restriction analysis and
15 sequence analysis.

For construction of a recombinant adenovirus containing sequences encoding the ATF.TIMP1 hybrid protein, this sequence is cloned in the pMAD5 expression plasmid. This plasmid contains part of the wildtype adenovirus type 5 DNA
20 sequences, a Major Late Promoter (MLP), and a polyadenylation (polyA) signal and can be used as either an expression vector or a shuttle vector to construct a recombinant adenovirus. To clone the ATF.TIMP1 sequence into the pMAD5 plasmid between the MLP promoter and the polyA
25 signal the following strategy has been followed.

Starting from a pCRII plasmid in which a 1373 base pair fragment of the uPA cDNA was cloned, a PCR reaction with the oligonucleotides 5'-cccggtctttttccatctgcgcagtc-3' and 5'-agggtcaccaaggaagagaatggc-3' was performed as described in
30 example 1 to make a pCRII-ATF plasmid (see figure 1). Subsequently on this pCRII-ATF plasmid a PCR reaction was performed using the oligonucleotides 5'-aatattattgaacttcatcaagttcc-3' and 5'-gactctagagcaaaaatgacaaccag-3' and the resulting DNA
35 fragment was cloned into the pCRII cloning vector. In this way the signal peptide of u-PA is removed and a SspI

restriction enzyme recognition site is introduced (underlined). The resulting plasmid DNA is designated pCRIIATF*.

In parallel a fragment of the cDNA of TIMP1 in pUC13-TIMP1 encoding amino acid residues -23 to 184 of the
5 TIMP-1 protein, including the signal peptide but lacking the stop codon, was amplified using the oligonucleotides
5'-agagagacaccagagaacccaccat-3' and
5'-aatattggctatctgggaccgcagg-3' containing a recognition site for the restriction enzyme SspI (underlined) and cloned into
10 a pCRII cloning vector. The resulting plasmid DNA is designated pCRII-TIMP1*.

This vector was subsequently digested with the restriction enzymes SspI and EcoRV and the TIMP1 containing DNA fragment was cloned into a EcoRV-SspI digested pCRII-ATF*
15 plasmid. The resulting plasmid containing the TIMP-ATF DNA fragment was called pCRII-TIMP-ATF. In a next step, the TIMP-ATF sequence was cloned into pMAD5. This was done by digestion of the pCRII-TIMP-ATF plasmid with the restriction enzymes EcoRV and SpeI, isolation of the TIMP-ATF encoding
20 DNA fragment and cloning of this fragment into the SnaBI and SpeI digested pMAD5 plasmid. The cloning was tested by restriction analysis and sequence analysis.

EXAMPLE 7

25 Vectors encoding hybrid proteins containing multiple copies of the BPTI unit coupled to the ATF domain have been constructed. To construct these multiple BPTI vectors, the following strategy is followed.

The pMAD5-ATF-BPTI described in EXAMPLE 4 is
30 digested with the restriction enzymes SspI and BstEII. In this way the vector is opened exactly in the open reading frame at the end of the BPTI sequence. The pCRII-BPTI plasmid described in EXAMPLE 2 is digested with NruI and BstEII resulting in a BPTI encoding DNA fragment with one blunt end
35 (NruI). The fragment was then monodirectionally cloned into the SspI BstEII pMAD5-ATF-BPTI vector. The thus constructed

plasmid named pMAD5-ATF-BPTI-BPTI was used as a shuttle vector for the construction of recombinant adenoviruses.

This approach can be repeated multiple times to construct vectors containing multiple BPTI-domains.

5

EXAMPLE 8

A vector encoding a hybrid protein containing both a BPTI unit and a TIMP1 unit coupled to the ATF domain has been constructed. To construct this BPTI-TIMP vector, the following strategy is followed.

The pMAD5-ATF-BPTI described in EXAMPLE 4 is digested with the restriction enzymes SspI and BstEII. In this way the vector is opened right behind the BPTI sequence. The pCRII-TIMP plasmid described in EXAMPLE 6 is digested with NruI and BstEII resulting in a TIMP1 encoding DNA fragment with one blunt end. The fragment was then cloned into the SspI BstEII pMAD5-ATF-BPTI vector. The thus constructed plasmid named pMAD5-ATF-BPTI-TIMP was used as a shuttle vector for the construction of recombinant adenoviruses.

20

EXAMPLE 9

To monitor the production of a functional ATF-BPTI hybrid protein after transfection of cells with pMAD5 or transduction with a recombinant replication-deficient ATF-BPTI encoding adenovirus, the following tests have been performed.

- The production of the hybrid ATF-BPTI protein by CHO cells transfected with the pMAD5-ATF-BPTI was analyzed using a uPA ELISA that recognizes the ATF, the aminoterminal fragment of u-PA. Production of ATF-BPTI was clearly detectable both after transient transfection of CHO cells with the pMAD5-ATF-BPTI plasmid (50-100 ng/ml/24hrs) and after transduction with an ATF-BPTI encoding adenoviral vector (up to 1.5 µg/ml/24hrs).

- The cell culture media of CHO cells transduced with the ATF-BPTI adenovirus were analyzed using western blotting

techniques. After electrophoresis and blotting, parallel filters were analyzed with polyclonal antibodies against either u-PA or BPTI (raised against Trasylol®). In both filters a signal was detected at the same expected position at approximately 20kDa. This indicates that the protein produced indeed contains fragments of u-PA and BPTI, thus that the hybrid protein is produced.

- The function as an inhibitor of plasmin activity of the ATF-BPTI protein was first analyzed in solution using dilutions of the culture medium of ATF-BPTI virus infected CHO cells (approximately 1.8 µg/ml). They were incubated with plasmin (1 nM) and the activity of plasmin was measured using a chromogenic substrate. Trasylol® dilutions were used as control references. Plasmin inhibition by ATF-BPTI medium was very effective, diluting the medium 1000x (i.e. 100 nM ATF-BPTI) resulted in a 50% inhibition of the activity of 1 nM plasmin, a similar inhibition as was observed with 100 nM Trasylol®. Thus the activity of ATF-BPTI is comparable to that of commercially available Trasylol® (Bayer, Germany).

- The function of ATF-BPTI as an inhibitor for plasmin bound to the cell surface via the interaction of the ATF domain with the u-PA receptor (uPAR) was tested using mouse cell lines that are either or not transfected with the human uPA receptor gene. These cells were incubated for 6 hrs with diluted medium of the ATF-BPTI virus-infected CHO cells. Cell extracts were made of the uPAR-transfected cells and the parental mouse cells lacking the human uPAR. Parallel cultures underwent a short acid treatment (pH 3, 3 min) before the cell extracts were made. This treatment will remove any u-PA or ATF bound to the u-PA receptor. The cell extracts were incubated with 1nM plasmin and the plasmin activity was determined. Plasmin activity could only be inhibited by the cell extract of the u-PAR containing cell line. No inhibition of plasmin activity was observed in the cell extracts of parental cell line, lacking the u-PA receptor, and in the acid-treated u-PAR containing cell line.

This clearly indicates that ATF-BPTI can function as a u-PAR bound plasmin inhibitor.

TABLE 1

	% INHIBITION OF PLASMIN ACTIVITY			
cell line	uPAR transfected cell line		parental cell line	
acid treatment	-	+	-	+
% inhibition	93%	0%	0%	0%

5

EXAMPLE 10

Cell-specific expression of ATF-BPTI in endothelial cells e.g. to specifically inhibit the migration of endothelial cells during angiogenesis, is achieved by cloning sequences of the promoter of the human pre-pro-endothelin 1 gene (nucleotide 2180-3680 of HUMEDN1B (GENBANK)) in front of the ATF-BPTI encoding DNA in an adenoviral vector. In this way, highly endothelial cell-specific expression of the ATF-BPTI hybrid protein can be obtained.

10

EXAMPLE 11

Proteolytic degradation of the extracellular matrix is a key event in many cell migration and tissue remodeling processes. This proteolytic matrix degradation is often found to be mediated by urokinase-type plasminogen activation. In order to test whether infection with an ATF-BPTI encoding adenovirus can inhibit plasmin mediated extracellular matrix degradation, an experiment was performed using human synoviocytes. These cells were infected with an ATF-BPTI adenovirus while they were seeded on an ³H-labeled extracellular matrix existing of bovine cartilage material. Profound inhibition of matrix degradation could be observed in the virus treated cells (figure 4) indicating that matrix

15

20

25

degradation can be inhibited by infecting cells with the ATF-BPTI encoding virus.

Figure 4 shows the degradation of cartilage matrix by human synoviocytes in the presence of plasminogen. Matrix is incubated with control medium (lane 1), synoviocytes (lane 2), synoviocytes infected with ATF-BPTI adenovirus (lane 3), and synoviocytes incubated with Trasylol® (100KIU/ml) (lane 4).

EXAMPLE 12

In the process of restenosis smooth muscle cell migration and vessel wall remodeling are key events in which plasmin mediated proteolysis of extracellular matrix components is involved. In vivo application of general plasmin inhibitors to interfere in this process may have systemic side effects. Application of a plasmin inhibitor to the surface of the migrating cells might prevent these side effects. Infection of the blood vessel wall with an ATF-BPTI adenovirus at a site where neointima formation can be expected, e.g. in a transplanted "coronary by-pass" graft, might be a ideal way to produce the ATF-BPTI locally, and thus inhibit plasmin activity in the direct surroundings of the migrating (smooth muscle) cells, resulting in a reduced neointima formation.

This principle was tested using human saphenous vein organ cultures, a model system in which neointima formation can be mimicked very realistically. In parallel cultures, either or not infected with an ATF-BPTI adenovirus, the neointima formation was analyzed after three and four weeks. In the untreated tissues a clear neointima formation could be observed. Profound inhibition of the neointima formation could be observed in the tissues treated with 10^{10} pfu/ml ATF-BPTI adenovirus.

Appendix

Description and Nucleotide sequence of the pMAD5-ATF-BPTI plasmid.

5

From	To	Description
1	184	adenovirus sequence 5'
10 184	447	adenovirus Major Late Promoter (MLP)
447	644	tripartite leader sequence (TPL)
685	1167	urokinase ATF sequence
1168	1353	bovine pancreas trypsin inhibitor sequence
1360	1443	urokinase 3' sequence (including stop codon)
15 1514	1615	sequence derived from pSP65 and LacZ
1616	1751	SV40 poly A signal
1752	7334	adenovirus sequence 3'
9831	8971	β -lactamase

20 Nucleotide sequence:

```

1      CATTTTCGCG GGAAACTGA ATAAGAGGAA GTGAAATCTG AATAATTTTG TGTTACTCAT
61     AGCGCGTAAT ATTTGTCTAG GGCCGCGGGG ACTTTGACCG TTTACGTGGA GACTCGCCCA
121    GGTGTTTTTC TCAGGTGTTT TCCGCGTTCC GGGTCAAAGT TGGCGTTTTA TTATTATAGT
25 181    CAGCTGATCG AGCGGTGTTT CGCGGTCTC CTCGTATAGA AACTCGGACC ACTCTGAGAC
241    GAAGGCTCGC GTCCAGGCCA GCACGAAGGA GGCTAAGTGG GAGGGGTAGC GGTGTTGTC
301    CACTAGGGGG TCCACTCGCT CCAGGGTGTG AAGACACATG TCGCCCTCTT CGGCATCAAG
361    GAAGGTGATT GGTTTATAGG TGTAGGCCAC GTGACCGGGT GTTCTGGAAG GGGGGCTATA
421    AAAGGGGGTG GGGGCGCGTT CGTCCTCACT CTCTTCCGCA TCGCTGTCTG CGAGGGCCAG
30 481    CTGTTGGGGC TCGCGTTGA GGACAACTC TTCGCGTCTT TTCCAGTACT CTTGGATCGG
541    AAACCCGTCG GCCTCCGAAC GGTACTCCGC CACCGAGGGA CCTGAGCGAG TCCGCATCGA
601    CCGGATCGGA AAACCTCTCG AGAAAGGCGT CTAACCAGTC GCTGATCGAT AAGCTAGCTT
661    ACGCGTACAT CTGCAGAATT CGGCTTAACT CTAGACCATG AGAGCCCTGC TGGCGCGCCT
721    GCTTCTCTGC GTCCTGGTCG TGAGCGACTC CAAAGGCAGC AATGAACTTC ATCAAGTTCC
35 781    ATCGAACTGT GACTGTCTAA ATGGAGGAAC ATGTGTGTCC AACAAGTACT TCTCCAACAT
841    TCACTGGTGC AACTGCCCAA AGAAATTCGG AGGGCAGCAC TGTGAAATAG ATAAGTCAAA

```

901 AACCTGCTAT GAGGGGAATG GTCACCTTTTA CCGAGGAAAG GCCAGCACTG ACACCATGGG
961 CCGGCCCTGC CTGCCCTGGA ACTCTGCCAC TGTCTTCAG CAAACGTACC ATGCCACAG
1021 ATCTGATGCT CTTAGCTGG GCCTGGGGAA ACATAATTAC TGCAGGAACC CAGACAACCG
1081 GAGGCGACCC TGGTGCTATG TGCAGGTGGG CCTAAAGCCG CTTGTCCAAG AGTGCATGGT
5 1141 GCATGACTGC GCAGATGGAA AAAAGCCCCG ACCTGACTTC TGCCTAGAGC CTCCATATAC
1201 GGGTCCCTGC AAGGCCAGAA TTATCAGATA CTTCTACAAC GCCAAGGCTG GGCTCTGCCA
1261 GACCTTTGTA TATGGCGGCT GCAGAGCTAA AAGAAACAAT TTCAAGAGCG CAGAGGACTG
1321 CATGAGGACC TGTGGTGGTA ATATTGGGCC CTGGGTCACC AAGGAAGAGA ATGGCTGGC
1381 CCTCTGAGGG TCCCCAGGGA GGAAACGGGC ACCACCCGCT TTCTTGCTGG TTGTCAATTT
10 1441 TGCTCTAGAG TCAAGCCGAA TTCTGCAGAT ATCGTCCATT CCGACAGCAT CGCCAGTCAC
1501 TATGGCGTGC TGCTAGAGGA TCCCCGGGCG AGCTCGAATT CCAGCTGAGC GCCGGTCGCT
1561 ACCATTACCA GTTGGTCTGG TGTCAAAAAT AATAATAACC GGGCAGGGGG GATTCTGAAC
1621 TTGTTTATTG CAGCTTATAA TGTTTACAAA TAAAGCAATA GCATCACAAA TTTCACAAAT
1681 AAAGCATTTT TTTCCTGCA TTCTAGTTGT GGTGTGTCCA AACTCATCAA TGTATCTTAT
15 1741 CATGTCTGGA TCTGGAAGGT GCTGAGGTAC GATGAGACCC GCACCAGGTG CAGACCCTGC
1801 GAGTGTGGCG GTAAACATAT TAGGAACCAG CCTGTGATGC TGGATGTGAC CGAGGAGCTG
1861 AGGCCCCATC ACTTGGTGCT GGCCTGCACC CGCGCTGAGT TTGGCTCTAG CGATGAAGAT
1921 ACAGATTGAG GTACTGAAAT GTGTGGGCGT GGCTTAAGGG TGGGAAAGAA TATATAAGGT
1981 GGGGGTCTTA TGTAGTTTGT TATCTGTTTT GCAGCAGCCG CCGCCGCCAT GAGCACCAAC
20 2041 TCGTTTGATG GAAGCATTGT GAGCTCATAT TTGACAACGC GCATGCCCCC ATGGGCCGGG
2101 GTGCGTCAGA ATGTGATGGG CTCCAGCATT GATGGTCGCC CCGTCTGCC CGCAAATCT
2161 ACTACCTTGA CCTACGAGAC CGTGTCTGGA ACGCCGTTGG AGACTGCAGC CTCCGCCGCC
2221 GCTTCAGCCG CTGCAGCCAC CGCCCGCGGG ATTGTGACTG ACTTTGCTTT CCTGAGCCCG
2281 CTTGCAAGCA GTGCAGCTTC CCGTTCATCC GCCCGCGATG ACAAGTTGAC GGCTCTTTTG
25 2341 GCACAATTGG ATTCTTTGAC CCGGGAACCT AATGTCGTTT CTCAGCAGCT GTTGGATCTG
2401 CGCCAGCAGG TTTCTGCCCT GAAGGCTTCC TCCCCTCCA ATGCGGTTTA AAACATAAAT
2461 AAAAAACCAG ACTCTGTTTG GATTTGGATC AAGCAAGTGT CTTGCTGTCT TTATTTAGGG
2521 GTTTTGCGCG CGCGGTAGGC CCGGACCAG CGGTCTCGGT CGTTGAGGGT CCTGTGTATT
2581 TTTTCCAGGA CGTGGTAAAG GTGACTCTGG ATGTTTCTAGT ACATGGGCAT AAGCCCGTCT
30 2641 CTGGGGTGGG GGTAGCACCA CTGCAGAGCT TCATGCTGCG GGGTGGTGTT GTAGATGATC
2701 CAGTCGTAGC AGGAGCGCTG GCGGTGGTGC CTAAAAATGT CTTTCAGTAG CAAGCTGATT
2761 GCCAGGGGCA GGCCCTGGT GTAAGTGTTC ACAAAGCGGT TAAGCTGGGA TGGGTGCATA
2821 CGTGGGGATA TGAGATGCAT CTTGGACTGT ATTTTATAGT TGGCTATGTT CCCAGCCATA
2881 TCCCTCCGGG GATTCATGTT GTGCAGAACC ACCAGCACAG TGTATCCGGT GCACTTGGGA
35 2941 AATTTGTCAT GTAGCTTAGA AGGAAATGCG TGGGAAGAACT TGGAGACGCC CTTGTGACCT
3001 CCAAGATTTT CCATGCATTC GTCCATAATG ATGGCAATGG GCCCACGGGC GGCGGCCTGG

3061 GCGAAGATAT TTCTGGGATC ACTAACGTCA TAGTTGTGTT CCAGGATGAG ATCGTCATAG
3121 GCCATTTTTA CAAAGCGCGG GCGGAGGGTG CCAGACTGCG GTATAATGGT TCCATCCGGC
3181 CCAGGGGCGT AGTTACCCCTC ACAGATTGTC ATTTCCACG CTTTGAGTTC AGATGGGGGG
3241 ATCATGTCTA CCTGCGGGGC GATGAAGAAA ACGGTTTCCG GGGTAGGGGA GATCAGCTGG
5 3301 GAAGAAAGCA GGTTCCTGAG CAGCTGCGAC TTACCGCAGC CCGTGGGCCC GTAAATCACA
3361 CCTATTACCG GGTGCAACTG GTAGTTAAGA GAGCTGCAGC TGCCGTCATC CCTGAGCAGG
3421 GGGGCCACTT CGTTAAGCAT GTCCCTGACT CGCATGTTTT CCCTGACCAA ATCCGCCAGA
3481 AGGCGCTCGC CGCCAGCGA TAGCAGTTCT TGCAAGGAAG CAAAGTTTTT CAACGGTTTG
3541 AGACCGTCCG CCGTAGGCAT GCTTTTGAGC GTTTGACCAA GCAGTTCCAG GCGGTCCAC
10 3601 AGCTCGGTCA CCTGCTCTAC GGCATCTCGA TCCAGCATAT CTCCTCGTTT CGCGGGTTGG
3661 GGCGGCTTTC GCTGTACGGC AGTAGTCGGT GCTCGTCCAG ACGGGCCAGG GTCATGTCTT
3721 TCCACGGGCG CAGGGTCCTC GTCAGCGTAG TCTGGGTCAC GGTGAAGGGG TGCGCTCCGG
3781 GCTGCGCGCT GGCCAGGGTG CGCTTGAGGC TGGTCTGCT GGTGCTGAAG CGCTGCCGGT
3841 CTTGCCCTTG CGCGTCGGCC AGGTAGCATT TGACCATGGT GTCATAGTCC AGCCCTCCG
15 3901 CGGCGTGGCC CTTGGCGCGC AGCTTGCCCT TGGAGGAGGC GCCGCACGAG GGGCAGTGCA
3961 GACTTTTGAG GGCGTAGAGC TTGGGCGCGA GAAATACCGA TTCCGGGGAG TAGGCATCCG
4021 CGCCGCAGGC CCCGCAGACG GTCTCGCATT CCACGAGCCA GGTGAGCTCT GGCCGTTCCG
4081 GGTCAAAAAC CAGGTTTCCC CCATGCTTTT TGATGCGTTT CTTACCTCTG GTTTCATGA
4141 GCCGGTGTCC ACGCTCGGTG ACGAAAAGGC TGTCCGTGTC CCCGTATACA GACTTGAGAG
20 4201 GCCTGTCTC GAGCGGTGTT CCGCGGTCTT CCTCGTATAG AAATCGGAC CACTCTGAGA
4261 CAAAGGCTCG CGTCCAGGCC AGCACGAAGG AGGCTAAGTG GGAGGGGTAG CGGTCGTTGT
4321 CCACTAGGGG GTCCACTCGC TCCAGGGTGT GAAGACACAT GTCGCCCTCT TCGGCATCAA
4381 GGAAGGTGAT TGGTTTGTAG GTGTAGGCCA CGTGACCGG TGTTCCTGAA GGGGGGCTAT
4441 AAAAGGGGGT GGGGGCGCGT TCGTCTCAC TCTCTCCGC ATCGCTGTCT GCGAGGGCCA
25 4501 GCTGTTGGGG TGAGTACTCC CTCTGAAAAG CGGGCATGAC TTCTGCGCTA AGATTGTCAG
4561 TTTCCAAAAA CGAGGAGGAT TTGATATCA CCTGGCCCGC GGTGATGCCT TTGAGGGTGG
4621 CCGCATCCAT CTGGTCAGAA AAGACAATCT TTTTGTGTC AAGCTTGGTG GCAAACGACC
4681 CGTAGAGGGC GTTGACAGC AACTTGCGA TGGAGCGCAG GGTTTGGTTT TTGTCGCGAT
4741 CGGCGCGCTC CTTGGCCGCG ATGTTTAGCT GCACGTATTC GCGCGCAACG CACCGCCATT
30 4801 CGGGAAAGAC GGTGGTGCGC TCGTCGGGCA CCAGGTGCAC GCGCCAACCG CGGTTGTGCA
4861 GGGTGACAAG GTCAACGCTG GTGGCTACCT CTCCGCGTAG GCGCTCGTTG GTCCAGCAGA
4921 GGCGGCCGCC CTTGCGCGAG CAGAATGGCG GTAGGGGGTC TAGCTGCGTC TCGTCCGGGG
4981 GGTCTGCGTC CACGGTAAAG ACCCCGGGCA GCAGGCGCGC GTCGAAGTAG TCTATCTTGC
5041 ATCCTTGCAA GTCTAGCGCC TGCTGCCATG CGCGGCGGC AAGCGCGCGC TCGTATGGGT
35 5101 TGAGTGGGGG ACCCATGGC ATGGGGTGGG TGAGCGCGGA GGCGTACATG CCGCAAATGT
5161 CGTAAACGTA GAGGGGCTCT CTGAGTATTC CAAGATATGT AGGGTAGCAT CTTCCACCGC

5221 GGATGCTGGC GCGCACGTAA TCGTATAGTT CGTGCGAGGG AGCGAGGAGG TCGGGACCGA
5281 GGTTGCTACG GGCGGGCTGC TCTGCTCGGA AGACTATCTG CCTGAAGATG GCATGTGAGT
5341 TGGATGATAT GGTGACGC TGAAGACGT TGAAGCTGGC GTCTGTGAGA CCTACCGCGT
5401 CACGCACGAA GGAGGCGTAG GAGTCGCGCA GCTTGTTGAC CAGCTCGGCG GTGACCTGCA
5 5461 CGTCTAGGGC GCAGTAGTCC AGGGTTTCCT TGATGATGTC ATACTTATCC TGTCCCTTTT
5521 TTTTCCACAG CTCGCGGTTG AGGACAACT CTTCGCGGTC TTCCAGTAC TCTTGATCG
5581 GAAACCCGTC GGCCTCCGAA CGGTAAGAGC CTAGCATGTA GAACTGGTTG ACGGCCTGGT
5641 AGGCGCAGCA TCCCTTTTCT ACGGGTAGCG CGTATGCCTG CGCGGCCTTC CGGAGCGAGG
5701 TGTGGGTGAG CGCAAAGGTG TCCCTGACCA TGACTTTGAG GTACTGGTAT TTGAAGTCAG
10 5761 TGTCGTCGCA TCCGCCCTGC TCCCAGAGCA AAAAGTCCGT GCGCTTTTTG GAACCGGGAT
5821 TTGGCAGGGC GAAGGTGACA TCGTTGAAGA GTATCTTTCC CGCGCGAGGC ATAAAGTTGC
5881 GTGTGATGCG GAAGGGTCCC GGCACCTCGG AACGGTTGTT AATTACCTGG GCGGCGAGCA
5941 CGATCTCGTC AAAGCCGTTG ATGTTGTGGC CCACAATGTA AAGTTCCAAG AAGCGCGGGA
6001 TGCCCTTGAT GGAAGGCAAT TTTTAAAGTT CCTCGTAGGT GAGCTCTTCA GGGGAGCTGA
15 6061 GCCCGTGCTC TGAAAGGGCC CAGTCTGCAA GATGAGGGTT GGAAGCGACG AATGAGCTCC
6121 ACAGGTCACG GGCCATTAGC ATTTGCAGGT GGTGCGGAAA GGTCTTAAAC TGGCGACCTA
6181 TGGCCATTTT TTCTGGGGTG ATGCAGTAGA AGGTAAGCGG GTCTTGTTCC CAGCGGTCCC
6241 ATCCAAGGTT CGCGGCTAGG TCTCGCGCGG CAGTCACTAG AGGCTCATCT CCGCCGA ACT
6301 TCATGACCAG CATGAAGGGC ACGAGCTGCT TCCCAAAGGC CCCCATCCAA GTATAGGTCT
20 6361 CTACATCGTA GGTGACAAAG AGACGCTCGG TCGGAGGATG CGAGCCGATC GGAAGA ACT
6421 GGATCTCCCG CCACCAATTG GAGGAGTGGC TATTGATGTG GTGAAAGTAG AAGTCCCTGC
6481 GACGGGCCGA AACTCTGTGC TGGCTTTTGT AAAAACGTGC GCAGTACTGG CAGCGGTGCA
6541 CGGGCTGTAC ATCCTGCACG AGGTTGACCT GACGACCGCG CACAAGGAAG CAGAGTGGGA
6601 ATTTGAGCCC CTCGCTGGC GGGTTTGGCT GGTGGTCTTC TACTTCGGCT GCTTGCTCTT
25 6661 GACCGTCTGG CTGCTCGAGG GGAGTTACGG TGATCGGAC CACCACGCCG CGCGAGCCCA
6721 AAGTCCAGAT GTCCGCGCGC GCGGTCGGA GCTTGATGAC AACATCGCGC AGATGGGAGC
6781 TGTCCATGGT CTGGAGCTCC CGCGGCGTCA GGTGAGGCGG GAGCTCCTGC AGGTTTACCT
6841 CGCATAGACG GGTGAGGCG CGGGCTAGAT CCAGGTGATA CCTAATTTC AGGGGCTGGT
6901 TGGTGGCGGC GTCGATGGCT TGCAAGAGGC CGCATCCCCG CGGCGCGACT ACGGTACCGC
30 6961 GCGGCGGGCG GTGGGCCGCG GGGGTGTCCT TGATGATGC ATCTAAAAGC GGTGACGCGG
7021 GCGAGCCCCC GGAGGTAGGG GGGGCTCCGG ACCCGCCGGG AGAGGGGGCA GGGGCACGTC
7081 GCGCGCCGCG GCGGGCAGGA GCTGGTGCTG CGCGCGTAGG TTCTGGCGA ACGCGACGAC
7141 GCGGCGGTTG ATCTCCTGAA TCTGGCGCCT CTGCGTGAAG ACGACGGGCC CGGTGAGCTT
7201 GAGCCTGAAA GAGAGTTCGA CAGAAATCAAT TTCGGTGTCG TTGACGGCGG CCTGGCGCAA
35 7261 AATCTCCTGC ACGTCTCCTG AGTTGTCTTG ATAGGCGATC TCGGCCATGA ACTGCTCGAT
7321 CTCTTCCTCC TGGAGATCAA TTGAAGCTAG CTTTAATGCG GTAGTTTATC ACAGTTAAAT

7381 TGCTAACGCA GTCAGGCACC GTGTATGAAA TCTAACAATG CGCTCATCGT CATCCTCGGC
7441 ACCGTACCCC TGGATGCTGT AGGCATAGGC TTGGTTATGC CGGTACTGCC GGGCCTCTTG
7501 CGGGATATCG TCCATTCCGA CAGCATCGCC AGTCACTATG GCGTGCTGCT AGCGCTATAT
7561 GCGTTGATGC AATTTCTATG CGCACCCGTT CTCGGAGCAC TGTCCGACCG CTTTGGCCGC
5 7621 CGCCCAGTCC TGCTCGCTTC GCTACTTGGA GCCACTATCG ACTACGCGAT CATGGCGACC
7681 ACACCCGTCC TGTGGATCTC GACCGATGCC CTTGAGAGCC TTCAACCCAG TCAGCTCCTT
7741 CCGGTGGGCG CGGGGCATGA CTATCGTCGC CGCACTTATG ACTGTCTTCT TTATCATGCA
7801 ACTCGTAGGA CAGGTGCCGG CAGCGCTCTG GGTCAATTTT GCGAGGACC GCTTTCGCTG
7861 GAGCGCGACG ATGATCGGCC TGTCGCTTGC GGTATTCGGA ATCTTGACG CCCTCGCTCA
10 7921 AGCCTTCGTC ACTGGTCCCG CCACCAAACG TTTCGGCGAG AAGCAGGCCA TTATCGCCCG
7981 CATGGCGGCC GACGCGCTGG GCTACGTCTT GCTGGCGTTC GCGACGCGAG GCTGGATGGC
8041 CTTCCCCATT ATGATTCTTC TCGCTTCCGG CGGCATCGGG ATGCCCGCGT TGCAGGCCAT
8101 GCTGTCCAGG CAGGTAGATG ACGACCATCA GGGACAGCTT CAAGGATCGC TCGCGGCTCT
8161 TACCAGCCCA GCAAAGGCC AGGAACCGTA AAAAGGCCGC GTTGCTGGCG TTTTTCATA
15 8221 GGCTCCGCCC CCCTGACGAG CATCACAAAA ATCGACGCTC AAGTCAGAGG TGGCGAAACC
8281 CGACAGGACT ATAAAGATAC CAGGCGTTTC CCCCTGGAAG CTCCCTCGTG CGCTCTCCTG
8341 TTCCGACCTT GCCGCTTACC GGATACCTGT CCGCCTTTCT CCCTTCGGGA AGCGTGGCGC
8401 TTTCTCATAG CTCACGCTGT AGGTATCTCA GTTCGGTGTA GGTGTTTCGC TCCAAGCTGG
8461 GCTGTGTGCA CGAACCCCCC GTTCAGCCCG ACCGCTGCGC CTTATCCGGT AACTATCGTC
20 8521 TTGAGTCCAA CCCGGTAAGA CACGACTTAT CGCCACTGGC AGCAGCCACT GGTAACAGGA
8581 TTAGCAGAGC GAGGTATGTA GCGGTGCTA CAGAGTTCTT GAAGTGTTGG CCTAACTACG
8641 GCTACACTAG AAGGACAGTA TTTGGTATCT GCGCTCTGCT GAAGCCAGTT ACCTTCGGAA
8701 AAAGAGTTGG TAGCTCTTGA TCCGGCAAAC AAACCACCGC TGGTAGCGGT GGTTTTTTTG
8761 TTTGCAAGCA GCAGATTACG CGCAGAAAAA AAGGATCTCA AGAAGATCCT TTGATCTTTT
25 8821 CTACGGGGTC TGACGCTCAG TGAACGAAA ACTCACGTTA AGGGATTTTG GTCATGAGAT
8881 TATCAAAAAG GATCTTCACC TAGATCCTTT TAAATTAAAA ATGAAGTTT AAATCAATCT
8941 AAAGTATATA TGAGTAAACT TGGTCTGACA GTTACCAATG CTTAATCAGT GAGGCACCTA
9001 TCTCAGCGAT CTGTCTATTT CGTTCATCCA TAGTTGCCTG ACTCCCCGTC GTGTAGATAA
9061 CTACGATACG GGAGGGCTTA CCATCTGGCC CAGTGCTGC AATGATACCG CGAGACCCAC
30 9121 GCTCACCGGC TCCAGATTTA TCAGCAATAA ACCAGCCAGC CGGAAGGGCC GAGCGCAGAA
9181 GTGGTCCTGC AACTTTATCC GCCTCCATCC AGTCTATTAA TTGTTGCCGG GAAGCTAGAG
9241 TAAGTAGTTC GCCAGTTAAT AGTTTGCACA ACGTTGTTGC CATTGCTGCA GGCATCGTGG
9301 TGTCACGCTC GTCGTTTGGT ATGGCTTCAT TCAGCTCCGG TTCCCAACGA TCAAGGCGAG
9361 TTACATGATC CCCATGTTG TGCAAAAAAG CGGTTAGCTC CTTGCGTCCT CCGATCGTTG
35 9421 TCAGAAGTAA GTTGGCCGCA GTGTTATCAC TCATGTTAT GGCAGCACTG CATAATTCTC
9481 TTA CTGTCAT GCCATCCGTA AGATGCTTTT CTGTGACTGG TGAGTACTCA ACCAAGTCAT

9541 TCTGAGAATA GTGTATGCGG CGACCGAGTT GCTCTTGCCC GCGTCAACA CGGGATAATA
9601 CCGCGCCACA TAGCAGAACT TTAAAAGTGC TCATCATTGG AAAACGTTCT TCGGGGCGAA
9661 AACTCTCAAG GATCTTACCG CTGTTGAGAT CCAGTTCGAT GTAACCCACT CGTGCACCCA
9721 ACTGATCTTC AGCATCTTTT ACTTTCACCA GCGTTTCTGG GTGAGCAAAA ACAGGAAGGC
5 9781 AAAATGCCGC AAAAAAGGGA ATAAGGGCGA CACGGAAATG TTGAATACTC ATACTCTTCC
9841 TTTTCAATA TTATTGAAGC ATTTATCAGG GTTATTGTCT CATGAGCGGA TACATATTG
9901 AATGTATTTA GAAAAATAAA CAAATAGGGG TTCCGCGCAC ATTTCCCCGA AAAGTGCCAC
9961 CTGACGTCTA AGAAACCATT ATTATCATGA CATTAACCTA TAAAAATAGG CGTATCACGA
10021 GGCCCTTTCG TCTTCAAGAA TTCTCATGTT TGACAGCTTA TCATCATCAA TAATATACCT
10 10081 TATTTTGGAT TGAAGCCAAT ATGATAATGA GGGGGTGGAG TTTGTGACGT GCGCGGGGGC
10141 GTGGGAACGG GCGGGGTGAC GTAGTAGTGT GCGGGAAGTG TGATGTTGCA AGTGTGGCGG
10201 AACACATGTA AGCGACGGAT GTGGCAAAAG TGACGTTTTT GGTGTGCGCC GGTGTACACA
10261 GGAAGTGACA ATTTTCGCGC GGTTTTAGGC GGATGTTGTA GTAAATTGG GCGTAACCGA
10321 GTAAGATTTG GC

Claims

1. A recombinant nucleic acid molecule comprising a vector useful for transfection or transduction of mammalian, e.g. human, cells, wherein said vector contains a nucleic acid insertion encoding an expressible hybrid polypeptide or protein which comprises a domain with a binding function and a domain with an effector function.
2. A recombinant nucleic acid molecule according to Claim 1, wherein said domain with a binding function comprises a receptor binding domain.
3. A recombinant nucleic acid molecule according to Claim 2, wherein said receptor binding domain is selected from the group consisting of urokinase receptor binding domain of urokinase, receptor binding domain of epidermal growth factor, receptor associated protein that binds to LDL Receptor related protein (α_2 -macroglobulin receptor) and VLDL Receptor.
4. A recombinant nucleic acid molecule according to Claim 2, wherein said receptor binding domain comprises the aminoterminal part of urokinase which is capable of binding to the urokinase receptor.
5. A recombinant nucleic acid molecule according to Claim 2, wherein said receptor binding domain comprises amino acid residues 1 through 135 of urokinase.
6. A recombinant nucleic acid molecule according to Claim 1, wherein said domain with an effector function is an enzymatically active domain.
7. A recombinant nucleic acid molecule according to Claim 1, wherein said domain with an effector function has protease inhibitor activity.
8. A recombinant nucleic acid molecule according to Claim 7, wherein said domain having protease inhibitor activity comprises a protease inhibitor or active part

thereof, said protease inhibitor being selected from the group consisting of (bovine) pancreatic trypsin inhibitor, (bovine) splenic trypsin inhibitor, urinary trypsin inhibitor, tissue inhibitor of matrix metalloproteinase 1,
5 tissue inhibitor of matrix metalloproteinase 2, tissue inhibitor of matrix metalloproteinase 3, and elastase inhibitor.

9. A recombinant nucleic acid molecule according to Claim 7, wherein said domain having protease inhibitor
10 activity comprises (amino acid residues 53 through 94 of) mature bovine pancreatic trypsin inhibitor.

10. A recombinant nucleic acid molecule according to Claim 7, wherein said domain having protease inhibitor activity comprises bovine splenic trypsin inhibitor.

15 11. A recombinant nucleic acid molecule according to Claim 7, wherein said domain having protease inhibitor activity comprises a tissue inhibitor of matrix metalloproteinases.

12. A recombinant nucleic acid molecule according to
20 Claim 1, wherein said domain with an effector function comprises (an active part of) two or more different protease inhibitors, or two or more copies of (an active part of) a protease inhibitor, or both.

13. A recombinant nucleic acid molecule according to
25 Claim 1, wherein said vector is selected from the group consisting of viral and non-viral vectors useful for transfection or transduction of mammalian cells.

14. A recombinant nucleic acid molecule according to Claim 1, wherein said vector is an adenovirus vector or a
30 retrovirus vector useful for transfection or transduction of human cells.

15. A recombinant nucleic acid molecule according to Claim 1, wherein said vector is an adenovirus vector based on shuttle vector pMAD5.

35 16. A recombinant nucleic acid molecule according to Claim 1, wherein said nucleic acid insertion encoding an

expressible hybrid polypeptide or protein is under the control of a cell- or tissue-specific promoter.

17. A recombinant nucleic acid molecule according to Claim 1, wherein said nucleic acid insertion encoding an
5 expressible hybrid polypeptide or protein is under the control of an endothelial cell-specific promoter, or a vascular smooth muscle cell-specific promoter, or a liver-specific promoter.

18. A process for preventing local proteolytic activity,
10 extracellular matrix degradation, cell migration, cell invasion, or tissue remodeling, comprising transfecting or transducing the cells involved or cells in their environment with a recombinant nucleic acid molecule as claimed in any one of the preceding Claims to obtain local expression of the
15 hybrid polypeptide or protein encoded by said nucleic acid molecule.

19. A process for producing a hybrid polypeptide or protein which comprises a domain with a binding function and a domain with an effector function, comprising transfecting
20 or transducing mammalian cells with a recombinant nucleic acid molecule as claimed in any one of Claims 1 to 17 to obtain expression of the hybrid polypeptide or protein encoded by said nucleic acid molecule, and optionally recovering the hybrid polypeptide or protein produced.

25




Fig. 1




Fig. 2




Fig. 3




Fig. 4

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 9/72, 15/62, C07K 14/81 // 19/00	A3	(11) International Publication Number: WO 98/51788 (43) International Publication Date: 19 November 1998 (19.11.98)
(21) International Application Number: PCT/NL98/00259 (22) International Filing Date: 11 May 1998 (11.05.98) (30) Priority Data: 97201423.7 12 May 1997 (12.05.97) EP (34) Countries for which the regional or international application was filed: NL et al. (71) Applicant (for all designated States except US): NEDERLANDSE ORGANISATIE VOOR TOEGEPAST-NATUURWETENSCHAPPELIJK ON- DERZOEK TNO [NL/NL]; Schoemakerstraat 97, NL-2628 VK Delft (NL). (72) Inventors; and (75) Inventors/Applicants (for US only): QUAX, Paulus, Hubertus, Andreas [NL/NL]; Jacob van Heemskercklaan 59, NL-2253 JX Voorschoten (NL). VERHEIJEN, Johan, Hendrikus [NL/NL]; Verdijkstraat 14, NL-2651 VB Berkel en Rodenrijs (NL). (74) Agent: SMULDERS, Th., A., H., J.; Verenigde Octrooibu- reaux, Nieuwe Parklaan 97, NL-2587 BN The Hague (NL).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> (88) Date of publication of the international search report: 20 May 1999 (20.05.99)
(54) Title: METHOD AND CONSTRUCT FOR INHIBITION OF CELL MIGRATION (57) Abstract A recombinant nucleic acid molecule comprising a vector useful for transfection or transduction of mammalian cells, wherein said vector contains a nucleic acid insertion encoding an expressible hybrid polypeptide or protein which comprises a domain with a binding function and a domain with an effector function. The domain with a binding function may comprise a receptor binding domain, and the domain with an effector function may have enzymatic activity, in particular protease inhibitor activity. The vector may be a viral (e.g. adenovirus or retrovirus) or non-viral vector useful for transfection or transduction of mammalian cells. The nucleic acid insertion encoding an expressible hybrid polypeptide or protein may be under the control of a cell- or tissue-specific promoter. A process for preventing local proteolytic activity, extracellular matrix degradation, cell migration, cell invasion, or tissue remodeling, comprising transfecting or transducing the cells involved or cells in their environment with the recombinant nucleic acid molecule to obtain local expression of the hybrid polypeptide or protein encoded thereby. A process for producing the hybrid polypeptide or protein by transfecting or transducing mammalian cells with the recombinant nucleic acid molecule to obtain expression and optionally recovering the hybrid polypeptide or protein produced.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

INTERNATIONAL SEARCH REPORT

International Application No

PCT/NL 98/00259

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N9/72 C12N15/62 C07K14/81 //C07K19/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 96 34009 A (RUTGERS THE STATE UNIVERSITY OF NEW JERSEY) 31 October 1996 see page 10, line 10 - page 20, line 2 ---	1,13-15, 18,19
X	WO 91 12328 A (FOWLKES DANA M ET AL) 22 August 1991 * the whole document, esp. page 1-2 and page 28, line 20 - page 39, line 26 * ---	1
X	WO 96 23814 A (CELL GENESYS INC) 8 August 1996 see page 9 - page 10 ---	1,18
X	EP 0 383 599 A (MERCK & CO INC) 22 August 1990 see the whole document --- -/--	1-3,6,19

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

19 February 1999

Date of mailing of the international search report

09.03.99

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

De Kok, A

INTERNATIONAL SEARCH REPORT

International Application No
PCT/NL 98/00259

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 28955 A (GLIEMANN JOERGEN ET AL) 2 November 1995 see the whole document ---	1,3
A	WO 95 11987 A (INCYTE PHARMA INC) 4 May 1995 see page 25, line 4 - page 27, line 13 see page 85, line 10 - page 87, line 17 ---	1-5,7, 18,19
A	WO 92 02553 A (DELTA BIOTECHNOLOGY LTD) 20 February 1992 cited in the application see page 3, line 10 - page 9, line 1 ---	1-5,7,8, 11,18
X	WO 95 17885 A (RUTGERS, THE STATE UNIVERSITY OF NEW JERSEY) 6 July 1995 see page 7, line 7 - line 10 see page 15, line 1 - page 23, line 36 see page 32, line 1 - page 34, line 6 ---	1,7,8
A	---	2-5
P,X	WO 97 25422 A (NISSIN FOOD PRODUCTS LTD) 17 July 1997 cited in the application see abstract ---	1-5,7
P,X	QUAX P H A ET AL: "Inhibition of neointima formation in cultured human saphenous vein segments by an adenovirus expressing an urokinase receptor binding plasmin inhibitor" CIRCULATION, vol. 96, no. 8-Suppl., 21 October 1997, page 1669 XP002084540 us see abstract nr.: 3741 ---	1-5, 13-15, 18,19
X	EP 0 439 954 A (SERAGEN INC) 7 August 1991 see page 1 - page 4 ---	1,6
X	WO 97 00949 A (MASSACHUSETTS INSTITUTE OF TECHNOLOGY) 9 January 1997 see page 6, line 27 - page 7, line 6 see page 12, line 26 - page 13, line 17 ---	1,6
X	US 5 504 001 A (FOSTER DONALD C) 2 April 1996 see column 3, line 5 - line 55 ---	1,6
A	WO 88 09344 A (CREATIVE BIOMOLECULES INC) 1 December 1988 see abstract ---	1,6
	-/--	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/NL 98/00259

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 95 13091 A (INTERNATIONAL TECHNOLOGY MANAGEMENT ASS) 18 May 1995 see abstract ----	1,6
A	EP 0 404 750 A (WASHINGTON UNIVERSITY) 27 December 1990 see abstract ----	8,12
A	WO 95 21601 A (PROTEIN ENGINEERING CORP.) 17 August 1995 see page 4, line 15 - page 5, line 24 ----	8,12
A	EP 0 623 676 A (AMGEN INC) 9 November 1994 see page 1 - page 4 -----	1,8,12

INTERNATIONAL SEARCH REPORT

International application No.
PCT/NL 98/00259

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.

3. ☒ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
1-15 18 19

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

This International Searching Authority found multiple (groups of) inventions in this international application, as follows:

1. Claims: 1-5, 13-15 completely and 18,19 partially

A vector useful for transfection of mammalian cells comprising a nucleic acid insertion encoding an expressible hybrid polypeptide which comprises a domain with a binding function and a domain with an effector function, said binding function comprising a receptor binding domain and said vector is selected from the group of viral and non-viral vectors especially an adenovirus or a retrovirus vector and its use.

2. Claims: 6-12 completely and 18, 19 partially

A vector useful for transfection of mammalian cells comprising a nucleic acid insertion encoding an expressible hybrid polypeptide which comprises a domain with a binding function and a domain with an effector function, said effector function comprising an enzymatically active domain or a protease inhibitor activity and its use

3. Claims: 16-17 completely and 18, 19 partially

A vector useful for transfection of mammalian cells comprising a nucleic acid insertion encoding an expressible hybrid polypeptide which comprises a domain with a binding function and a domain with an effector function, wherein said nucleic acid insertion is under the control of a cell- or tissue-specific promoter and its use

INTERNATIONAL SEARCH REPORT

Information on patent family members

In International Application No

PCT/NL 98/00259

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9634009 A	31-10-1996	US 5843724 A AU 5676096 A	01-12-1998 18-11-1998
WO 9112328 A	22-08-1991	CA 2075974 A EP 0515516 A JP 5503430 T US 5498538 A US 5747334 A US 5625033 A US 5852167 A US 5844076 A	16-08-1991 02-12-1992 10-06-1993 12-03-1996 05-05-1998 29-04-1997 22-12-1998 01-12-1998
WO 9623814 A	08-08-1996	US 5712149 A AU 4861396 A CA 2221629 A EP 0842194 A US 5686281 A	27-01-1998 21-08-1996 08-08-1996 20-05-1998 11-11-1997
EP 0383599 A	22-08-1990	AT 133993 T AU 617039 B AU 4994690 A AU 631200 B CA 2010256 A CY 1987 A DE 69025211 D DE 69025211 T DK 383599 T ES 2085329 T FI 102383 B GR 3019836 T HK 15597 A IE 71934 B JP 2069035 C JP 2276590 A JP 7106159 B JP 2510846 B JP 7278200 A NO 300595 B PT 93178 A, B	15-02-1996 14-11-1991 23-08-1990 19-11-1992 17-08-1990 05-09-1997 21-03-1996 19-09-1996 05-08-1996 01-06-1996 30-11-1998 31-08-1996 14-02-1997 12-03-1997 10-07-1996 13-11-1990 15-11-1995 26-06-1996 24-10-1995 23-06-1997 31-08-1990
WO 9528955 A	02-11-1995	AU 2342995 A	16-11-1995
WO 9511987 A	04-05-1995	AU 685187 B AU 8076994 A EP 0730660 A JP 9504174 T US 5766897 A	15-01-1998 22-05-1995 11-09-1996 28-04-1997 16-06-1998
WO 9202553 A	20-02-1992	GB 2246779 A AU 8318591 A	12-02-1992 02-03-1992
WO 9517885 A	06-07-1995	US 5550213 A AU 1515295 A	27-08-1996 17-07-1995
WO 9725422 A	17-07-1997	AU 1210797 A EP 0890638 A	01-08-1997 13-01-1999
EP 0439954 A	07-08-1991	AU 657087 B	02-03-1995

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/NL 98/00259

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0439954	A		AU 7168991 A	24-07-1991
			AU 8032194 A	27-04-1995
			CA 2071969 A	23-06-1991
			JP 5502880 T	20-05-1993
			WO 9109871 A	11-07-1991
			US 5668255 A	16-09-1997
WO 9700949	A	09-01-1997	US 5726050 A	10-03-1998
US 5504001	A	02-04-1996	NONE	
WO 8809344	A	01-12-1988	AT 120761 T	15-04-1995
			AU 612370 B	11-07-1991
			AU 1804988 A	21-12-1988
			AU 648591 B	28-04-1994
			AU 8579991 A	13-02-1992
			DE 3853515 D	11-05-1995
			DE 3853515 T	17-08-1995
			EP 0318554 A	07-06-1989
			EP 0623679 A	09-11-1994
			JP 2500329 T	08-02-1990
			US 5482858 A	09-01-1996
			US 5476786 A	19-12-1995
			US 5132405 A	21-07-1992
			US 5091513 A	25-02-1992
			US 5258498 A	02-11-1993
WO 9513091	A	18-05-1995	AU 1176595 A	29-05-1995
EP 0404750	A	27-12-1990	AT 109796 T	15-08-1994
			DE 69011433 D	15-09-1994
			DK 404750 T	03-10-1994
			ES 2060134 T	16-11-1994
WO 9521601	A	17-08-1995	CA 2180950 A	17-08-1995
			EP 0739355 A	30-10-1996
			JP 9511131 T	11-11-1997
			US 5795865 A	18-08-1998
EP 0623676	A	09-11-1994	AT 170560 T	15-09-1998
			AU 648505 B	28-04-1994
			AU 5638890 A	18-12-1990
			CA 2017166 A	19-11-1990
			DE 69032609 D	08-10-1998
			EP 0398753 A	22-11-1990
			ES 2123495 T	16-01-1999
			JP 4500683 T	06-02-1992
			SG 48766 A	18-05-1998
			SG 43824 A	14-11-1997
			WO 9014363 A	29-11-1990
			US 5714465 A	03-02-1998